

**33 Congreso Argentino de Producción Animal
Comarca Viedma, Patagonia – 13 al 15 octubre 2010
Seccion Sistemas de Produccion**

Global Climate Change and Livestock Production

**Dr. Horst Jürgen Schwartz
Professor (retired)
Chair for Livestock Ecology
Humboldt University of Berlin**

Global Climate Change and Livestock production: Relevance, Consequences & Perspectives


Why do we keep livestock?

**Does livestock production contribute to
climate change?**

Is livestock production a victim of climate change?

Main functions of livestock production for society

Food security



Main functions of livestock production for society

Food security

Services



Main functions of livestock production for society

Food security

Services

Income generation



Numbers and biomass of domestic animals and humans

Species	Numbers (million)	Biomass (million t)
Cattle & Buffalo	1 480	540
Sheep	1 065	40
Goats	780	27
Equines	118	42
Camelids	24	8.5
Pigs	936	122
Poultry	14 711	15
Total		794.5
Humans	6 800	374

Main functions of the environment (nature) for livestock production

Production base



Main functions of the environment (nature) for livestock production

Production base

**Production
reserve**



Main functions of the environment (nature) for livestock production

Production base

**Production
reserve**

**Emission
depository**



Global Climate Change

**Is the contribution by “ruminant”
livestock substantial?**

YES

The ruminant as a production system

Inputs

large quantity of low nutrient density
plant materials
&
water

**intake by
ruminant**

RUMEN

anaerobic fermentation


animal metabolism
reproduction
growth
activity

**Heat losses
Material losses
Indirect losses**

**High nutrient density foods
Non-food products
Services**

Outputs

small
quantity



Climate affecting emissions from livestock production

Gaseous emissions


- CO_2 , carbon-dioxide
- CH_4 , methane
- NH_3 , ammonia & nitrous oxides
- Sulphur compounds

Dust

Ash

Soot particles

Proportion of various sources in the global methane emission



Source: Walker, 1994

**Climate affecting emissions out of
livestock originate from:**



**Primary production:
Carbon-dioxide, Methane, Nitrous Oxide**




**Secondary Production:
Carbon-dioxide, Methane, Ammonia,
Sulphur Compounds, Dust**

Annual methane losses from a model livestock production system: dairy farming in S.W. England


(102 cows, 110 others, stall feeding of silage and concentrate)

type of loss	total emission kg CH₄ - C year⁻¹
losses from ruminants	6775
losses from stored wastes	2285
losses from silage effluent	2596
losses from dirty water	332
total losses	11988



**Cultivation of Virgin Land:
Carbon Dioxide, Methane, Nitrous Oxides,
Ozone, Ash, Black Soot**



Agricultural expansion in Brazil



Dependence on agricultural exports

Beef exports are fastest growing commodity


Herd growth concentrated in Central and Northern Brazil



Deforestation:
 20000 km² • yr⁻¹
 50 to 60% for
 cattle ranching


■ Rainforest ■ Cerrado ■ Deforestation

Source: D. Bungenstab, 2004




**Burning of Agricultural Biomass:
Carbon Dioxide, Methane, Nitrous Oxides,
Ozone, Ash, Black Soot**

Relative contribution of biomass burning to various climate affecting emissions [% of all emissions]



Source: Levine et al.; 1995

Contribution of burning various types biomass and the resulting release of carbon into the atmosphere




Source: Levine et al.; 1995

Local machine times and agricultural transports




Transport routes of importance for Western European dairy production




Regional distribution of **CATTLE** populations and proportion of regional production of meat and milk

Source: FAO Production Yearbook 2004


Populations



Milk



Meat






**Black-and-white dairy cattle
yielding from 6000 to 10000 litres
of milk in a lactation of 9 months**



Holstein-Friesian

Annual net trade ['000 t] of feeds and dry milk powder between the major economic regions of the world from 1970 to 2000



Source: FAO Trade Yearbooks 1970-2000

Enteric fermentation CH₄ emission levels of ruminants are affected by:

- **animal species and genotype**
- **animal age and nutritional status**
- **animal performance level**
- **feed availability and intake level**
- **feed quality**
 - nutrient density, digestibility, protein-energy ratio**
 - seasonal variation**
 - botanical composition**
 - pasture and range management**

Enteric fermentation CH₄ emission for dairy cattle in relation to average milk yield per COW

Region	Kg CH ₄ /head/year	Kg Milk/head/year	Kg Milk/Kg CH ₄
North America	118	6700	57
Western Europe	100	4200	42
Eastern Europe	81	2550	32
Oceania	68	1700	25
Latin America	57	800	14
Asia	56	1650	29
Indian Subcontinent	46	900	20
Africa and Middle East	36	475	13

Source: IPCC Guidelines for National Greenhouse Gas Inventories and Authors' Calculations

Manure management systems affecting loss of CH₄, N₂O, trace gases

Liquid systems

- with storage lagoons
- with storage pits or silos
- with forced drying and grinding
- with biogas production



Manure management systems affecting loss of CH₄, N₂O, trace gases

Liquid systems

- with storage lagoons
- with storage pits or silos
- with forced drying and grinding
- with biogas production



Manure management systems affecting loss of CH₄, N₂O, trace gases

Liquid systems

- with storage lagoons
- with storage pits or silos
- with forced drying and grinding
- with biogas production



Manure management systems affecting loss of CH₄, N₂O, trace gases

Solid systems


- with or without bedding, dry stockpiling
- with or without bedding, composting
- with fuel use



Manure management systems affecting loss of CH₄, N₂O, trace gases

Pastoral systems

- with partial collection for fuel use
- with complete spreading by grazing stock



Manure management systems affecting loss of CH₄, N₂O, trace gases

Liquid systems

- with storage lagoons
- with storage pits or silos
- with forced drying and grinding
- with biogas production


Solid systems

- with or without bedding, dry stockpiling
- with or without bedding, composting
- with fuel use

Pastoral systems

- with partial collection for fuel use
- with complete spreading by grazing stock

losses




Manure management CH₄ emission factors [kg CH₄/head/year] for dairy cattle in relation to region and climate type

Region	Cool	Temperate	Warm
North America	36	54	76
Western Europe	14	44	81
Eastern Europe	6	19	33
Oceania	31	32	33
Latin America	0	1	2
Asia	7	16	27
Africa and Middle East	1	1	2
Indian Subcontinent	5	5	6

Source: IPCC Guidelines for National Greenhouse
Gas Inventories

**Worldwide distribution of crops ■ and pastures ■
as the base for livestock production systems**



POSSIBLE TOOLS FOR THE CONTROL OF AGRICULTURAL EMISSIONS

Technical emission controls

- storing and handling
- application to land
- reduction of N-leaching
- mechanical treatment

Biological emission controls

- feeding the animals
- biological treatment of feeds
- manipulation of rumen microbes

Economic emission controls

- emission taxes
- consumption taxes
- product taxes
- subsidies for clean production
- emission quotas
- transferable emission quotas

Legal emission controls